

## A new species of *Pingasa* (Lepidoptera, Geometridae, Geometrinae) from Hong Kong

A. C. GALSWORTHY

11 Church Path, Merton Park, London SW193HJ, UK

**Abstract** A new species of *Pingasa* is described from Hong Kong and SE Asia.

**Key words** *Pingasa*, new species, Hong Kong, China, taxonomy.

### Introduction

Holloway (1982) noted the existence in Hong Kong of an undescribed species of *Pingasa*. During recent work in Hong Kong further specimens have been taken. It is now named as :

### *Pingasa chloroides* sp. nov.

Male (Fig. 1). Forewing 19 mm ; very similar in facies to *Pingasa chlora* Stoll. The crenellated postmedial on the hindwing is bordered on the outer side with pale brown (white in *chlora*). The white markings in the submarginal area are generally less extensive in *chloroides* sp. nov., and the white area basal to the postmedial has a more uniform dusting of pale brown scales. The forewing antemedial line is subtly different, the crenellations being more sinuous, without the prominent acute angle in *chlora*.

Female. Similar to the male.

Male genitalia (Fig. 2). Differ principally from congeners in having the tips of the valves deeply bifid, each lobe terminating in a sharp point, the whole resembling a lobster claw.

Holotype. ♂, Lam Tsuen Valley, Hong Kong, 8. iii. 1997, R. Kendrick, BMNH, BM




Fig. 1. *Pingasa chloroides* sp. nov., male.




Fig. 2. *Pingasa chloroides* sp. nov., male genitalia.

Geometrid slide no 19189. Paratypes: 1 ♂, Victoria Peak, Hong Kong, 22. vi. 1922, coll A. C. Galsworthy; 1 ♂, Lai Chi Chong, Hong Kong, 12. vi. 1997, BMNH; 1 ♂, Yuen Long, Hong Kong, 19. xi. 1987, K. Li, BMNH; 1 ♀, Fauling, SE China, ix. 1911 (E. Wahr), Rothschild collection, BM 1939-1; 1 ♂, Cochin China, Nacham, vii. 1936, S. Masseyeff, BM 1937-658; 1 ♂, Hong Kong, ii. 1892, JOW, BM 92-114, BM Geometrid slide no 11061.

Rasge. SE China, Hong Kong, Indo-China.

Remarks. This species resembles *P. secreta* Inoue from Taiwan, but differs in narrower forewings and more elongated hindwing, narrower brown border on both wings, crenellated postmedial on hindwing straighter (curved towards the inner margin in *secreta*) and no or very little white on outer edge of postmedial of hindwing, and in male genitalia. There is a single female specimen from Taiwan (Formosa, Takow, 24. v. 1905, A. E. Wileman, BM 1913-180) in the BMNH which appears to belong to this species, but in the absence of male material from Taiwan it has been omitted from the type series.

### Acknowledgements

I am grateful to the Trustees and staff of the Natural History Museum, London, for allowing me access to the collection and libraries. I should also like to thank Mr Kent Li, Mr R. Kendrick and Mr G. Reels, together with the Department of Ecology and Biodiversity of the University of Hong Kong, for supplying material, and Dr J. D. Holloway and an anonymous referee for commenting on this paper.

### Reference

- Holloway, J. D. 1982. Taxonomic appendix. In Barlow, H. S., *An Introduction to the Moths of South East Asia*: 174-271. Kuala Lumpur.

## 摘 要

香港産 *Pingasa* の 1 新種 (A. C. Galsworthy)

香港から *Pingasa* 属のオアシャクの 1 新種, *P. chloroides* sp. nov. を記載した. 外観は *P. chlora* Stoll に似るが, ♂交尾器 valva は先端で深く二分し, エビの鉗のように鋭く尖る点で, 他の同属種とも区別できる. 本新種は, 香港の他, 中国南東部, インドシナにも分布する.

[文責: 吉本 浩]

(Accepted November 21, 1997)